

# Cryptosporidium Transport in Unsaturated Flow

Christophe Darnault, Tammo Steenhuis  
J.-Yves Parlange, Philippe Baveye, David Cornwell

USEPA/USGS Meeting – Cryptosporidium Removal by Bank Filtration  
September 10, 2003 – Reston, Virginia



Cornell University  
Dept. of Biological and  
Environmental Engineering




ENVIRONMENTAL ENGINEERING & TECHNOLOGY, INC.


## Structure of Seminar

- Preferential Flow and Vadose Zone
- Preferential transport of solute/biocolloid *Cryptosporidium parvum* oocysts through vadose zone
- Take-home Messages

# Vadose Zone



# Preferential Flow




- Process
  - By-pass transport phenomena
  - Concentration of flow into channels at soil surface or subsurface
- Types
  - Macropore flow
  - Fingered flow
  - Funneled flow
- Impacts on water resources
  - Non-point sources pollution
  - Water quality

# Preferential Transport of *Cryptosporidium*

- Characterize the fate and transport of *Cryptosporidium* in the subsurface environment
- Modeling of solute and biocolloid transport


## Issues




- *Cryptosporidium parvum*
  - 4-6  $\mu\text{m}$  diameter
  - Found in feces of infected animals
  - Protozoan pathogen that causes Cryptosporidiosis
- Source of *Cryptosporidium parvum*
  - Application of manure to farm fields
  - Wild animals
- *Cryptosporidium* in drinking water
  - Resistant to chlorination

# Laboratory Experiments

- Laboratory column experiments
  - *Cryptosporidium parvum* from calves feces
  - Rainfall simulation event
  - Different porous media: silica sand, water repellent sand, sand with water repellent layers
  - Undisturbed soil columns
  - Mixture of *Cryptosporidium parvum* and a tracer (chloride)



# Microbiology Analysis



Dye-uptake types

(Anguish L.J. and W.C. Ghiorse. 1997)



- Oocysts Visualization & Viability
  - Immunofluorescence Staining
  - Dye Permeability Assay
- Soil extraction protocols
  - Tween & gradient concentration by centrifugation
- Fluorescence Microscopy
  - Nonviable oocysts:
    - permeable to DAPI and PI
  - Viable oocysts:
    - impermeable to DAPI and PI, or permeable to DAPI only

# Results

- Visualization of preferential flow path
- Visualization, quantification of oocysts
- Breakthrough curves (BTC)
- *Cryptosporidium* distribution and water saturation in soil profile
- Modeling of solute and colloid transport

# Visualization of Fingering Flow



## Water Fingering Flow in Sand



3D - 2D Visualization & Fluid Content Profile



# BTC of Cl and Oocysts & Soil Profile Distribution of Water and Oocysts

1 cm/hr rainfall, 12/20 sand





# BTC of Cl and Oocysts & Soil Profile Distribution of Water and Oocysts

2 cm/hr rainfall, 12/20 sand




# BTC of Cl and Oocysts & Soil Profile Distribution of Water and Oocysts


1 cm/hr rainfall, 12/20 sand  
with two water repellent interfaces layers





# Undisturbed Soil Column



# BTC of Cl and Oocysts



# Schematic and Model for Preferential Flow



Distribution Layer (DL)  
Exponential loss of solutes/colloids

$$C = C_0 \exp(-\lambda t)$$

Conveyance Zone (CZ)  
Convective-dispersive equation

$$\frac{\partial C}{\partial t} = D \frac{\partial^2 C}{\partial x^2} - v \frac{\partial C}{\partial t}$$


# Observed & Predicted BTC

Solutes concentration

$$C = \frac{1}{2} C_0 \exp(-\lambda t) \left[ \exp\left\{ \frac{vx}{2D} (1-\alpha) \right\} \operatorname{erfc}\left( \frac{x-vt\alpha}{2\sqrt{Dt}} \right) + \exp\left\{ \frac{vx}{2D} (1+\alpha) \right\} \operatorname{erfc}\left( \frac{x+vt\alpha}{2\sqrt{Dt}} \right) \right]$$

Colloids concentration

$$C = \frac{1}{2} C_0 \exp\left[ -\left( \frac{q}{W} + \beta \right) t \right] \left[ \exp\left[ \frac{vx}{2D} (1-\alpha') \right] \operatorname{erfc}\left( \frac{x-vt\alpha'}{\sqrt{4Dt}} \right) \right]$$



# Unsaturated Zone and Bank Filtration

- Unsaturated zone and its occurrence in bank filtration
  - well production over-pumping
  - during high river stage
  - flooding of dry river bench
- Unsaturated zone and its role in fate and transport of contaminants
  - preferential transport phenomena
  - gas-water interfaces
  - physical, chemical and biological processes
  - contaminants attenuation and entrapment

# Take-home Messages

- Demonstrated fast transport of *Cryptosporidium parvum* oocysts by fingered and macropores flow through vadose zone
- Experiments results suggest that human pathogens, like *Cryptosporidium parvum* oocysts, can be rapidly transported to significant depths in situations where preferential flow occurs
- Gas-water interfaces limit the movement of oocysts
- Modeled preferential transport of solutes and colloids
- Unsaturated zone and its importance in bank filtration

# Acknowledgements

Cornell University, USA  
U. S. Department of Agriculture